

# Diagenetic origin for quartz-pebble conglomerates

Rónadh Cox\*

Ethan D. Gutmann\*

Patricia G. Hines

Geosciences Department, Williams College, Williamstown, Massachusetts 01267, USA

## ABSTRACT

**The occurrence of quartz-pebble conglomerates (QPC) in the rock record increases backward through time from the Tertiary through the Precambrian. The positive correlation between QPC abundance and age is valid both for numbers of reported QPC and for QPC as a percentage of all conglomerate, and at both the era and the period level. QPC are usually interpreted as being due to intense chemical weathering, protracted transport, or sediment recycling, but none of these can account for the age distribution of QPC, which is the opposite of the global mass-age distribution for sedimentary rocks. Precambrian and Tertiary conglomerates with similar sources and sedimentology have vastly different clast populations, nonquartzose clasts being much more abundant in the younger rocks. Comparison of the petrology of QPC and polymict conglomerates shows that QPC have consistently higher proportions of diagenetic secondary matrix and pressure-solved grain contacts. We conclude that diagenetic factors play an important role in QPC formation by preferentially destroying less durable clasts.**

**Keywords:** clasts, diagenesis, pressure solution, quartz-pebble conglomerates.

## INTRODUCTION

In quartz-pebble conglomerates (QPC), more than 90% of the clasts consist of vein quartz, chert, or quartzite (Boggs, 1992). Some ancient QPC are >100 m thick and cover areas ~25 km<sup>2</sup> (Trevena, 1979; Ethridge et al., 1984; Mosher et al., 1993; Cherichetti et al., 1998), and interbedded sandstone-QPC sequences can be >1000 m thick and cover hundreds of square kilometers (Kingsley, 1984), but no volumetrically significant deposits of this type are forming at present. Small quartz-pebble accumulations may occur locally in drainages directly overlying quartzite bedrock, but thick or laterally extensive modern units are not known. In the geologic record, however, substantial QPC accumulations are found in sequences derived from lithologically diverse source rocks. In addition, QPC become more common further back in the geologic record.

QPC are generally considered to have specific paleoclimatic or paleogeographic implications. Current sedimentology textbooks state that they represent tectonically quiescent conditions under which chemical and mechanical weathering were very efficient (e.g., Prothero and Schwab, 1996, p. 76; Selley, 2000, p. 383; Boggs, 2001, p. 151). Processes invoked to explain QPC include prolonged mechanical abrasion (Abbott and Peterson, 1978; Kingsley, 1984), intense chemical weathering (Dal Cin, 1968; Reimer and Mossman, 1990), or recycling of older conglomerate (Youngson and Craw, 1996). Many QPC,

however, were deposited on alluvial fans, implying a short interval between erosion and deposition and thus little time for breakdown of labile clasts. In addition, independent evidence for intense weathering conditions at source or in the depositional basin is generally lacking. There are abundant examples of such problematic deposits (e.g., Smith, 1967; Ethridge et al., 1984; Kingsley, 1984; Kraus, 1984; Bayne, 1987; Mosher et al., 1993).

QPC host many gold and uranium orebodies, and there is active debate about connections between the mineralization and diagenesis (e.g., Robinson and Spooner, 1984; Reimer and Mossman, 1990). Conglomerates also form petroleum reservoirs (e.g., Glover, 1982; Cronin and Kidd, 1998), and prediction of porosity, permeability, and reservoir quality depend on understanding their diagenetic histories. More generally, understanding QPC genesis bears on interpretations of paleogeography, paleoclimate, and depositional environments. We present evidence that QPC may form during diagenesis by alteration of polymict conglomerate precursors.

## AGE DISTRIBUTION OF QUARTZ-PEBBLE CONGLOMERATES

Sedimentary rocks show a well-documented inverse relationship between preserved volume and age (Gilluly, 1969; Garrels et al., 1972; Ronov et al., 1980; Ronov, 1983). QPC, in contrast, tend to be more common in older sequences. It is not possible to measure the global mass-age distribution of conglomerates directly, because the units are generally below the scale of regional geologic maps. We have therefore used the American Geological Insti-

tute GeoRef database as a proxy (Appendix<sup>1</sup>) and have tallied references to conglomerates and QPC of different ages (Table 1; see footnote 1).

Conglomerate records (QPC plus non-QPC) constitute 2%–3% of the GeoRef records for each time interval from Tertiary to Archean, reflecting the fact that conglomerates form a minor but consistent proportion of sedimentary rocks (Table 1, column B; see footnote 1). In addition, the records for conglomerates (QPC plus non-QPC) show a marked decrease with increasing age, which closely matches the global mass-age distribution of sedimentary rocks (Fig. 1). QPC data, however, have a very different age distribution. The number of records for Paleozoic QPC is double that for the Cenozoic and Mesozoic, and for the Precambrian it increases further, by a factor of three (Fig. 2). The trend does not reflect variation in conglomerate volume through time, because the QPC numbers change as a proportion of total conglomerate records (Table 1, column C; see footnote 1). Likewise, the consistency of the proportion of conglomerate records for each time interval, including the Proterozoic and Archean (Table 1, column B; see footnote 1), demonstrates that the pattern is not an artifact of era or period length.


QPC are most prominent in the Precambrian. Whereas average Phanerozoic values are <1%, QPC account for 6.4% of Proterozoic and 6% of Archean conglomerate records. The database for Precambrian QPC of known age is small and the time intervals are large in comparison with the Phanerozoic (Table 1, column D; see footnote 1), but the pattern is the same: the number and proportion of QPC increase strongly with increasing age. This consistent trend toward increasing abundance of QPC in older rocks is in direct contrast to the age distribution of preserved sedimentary rock, and requires explanation.

## DIAGENETIC EFFECTS ON THE COMPOSITION OF CLASTIC ROCKS

Diagenesis plays a major role in the compositional evolution of clastic rocks. Chemical alteration coupled with mass transfer of ma-



\*E-mail: Cox—rcox@Williams.edu. Present address: Gutmann—University of Colorado, Boulder, Colorado 80309, USA.

<sup>1</sup>Data Repository item 2002031, Appendix, Description of the search terms used and the rationale, Table 1, GeoRef data for conglomerates and QPC, and Table 2, Point-count data for thin sections of conglomerate interclast material, is available from Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301-9140, editing@geosociety.org, or at www.geosociety.org/pubs/ft2002.htm.



**Figure 1.** Comparison between numbers of GeoRef records for conglomerates (both quartz-pebble conglomerates [QPC] and non-QPC) and mass-age distribution of sedimentary rocks. Survival rate for each time interval (axis A) is given in metric tons per year (graph shows data from Gregor, 1985; other models are very similar).

material can produce extensive secondary porosity (Siebert et al., 1984; Surdam et al., 1984) and ultimately diagenetic quartz arenites (e.g., Milliken, 1988; Abdel Wahab, 1998). Most studies have focused on sandstones, but the principles should be equally applicable to conglomerates.



**Figure 2.** Quartz-pebble conglomerate (QPC) records for each time interval as percentage of all conglomerate (QPC plus non-QPC) records (see also Table 1 [text footnote 1]).

Volume loss, mostly through dissolution of feldspar and rock fragments, is commonly 20%–30% (Wilkinson et al., 1997; Abdel Wahab, 1998). Intergranular volumes <30% are commonly seen in well-sorted sandstones, indicating dissolution and loss of grain mass (Houseknecht, 1987, 1989). There are marked compositional gradients with depth due to intrastratal dissolution (Cavazza and Gandolfi, 1992), and volume loss of almost 40% is known in deeply buried sandstone (Milliken et al., 1994). In the absence of overpressuring, secondary porosity can be eliminated by compaction (Harris, 1989), in which case there may be no record whatsoever of the former presence of a labile grain population.

To our knowledge, there are no studies of porosity loss, diagenetic mass transfer, or intrastratal solution in conglomerates. Pressure

solution in QPC is very common, however, at both the macroscopic and microscopic levels (Mosher, 1976, 1981). Diagenetic quartz arenites have also been shown to display an anomalously high proportion of strongly welded and pressure-solved grain contacts (Harris, 1989).


### DIAGENESIS IN POLYMICT AND QUARTZ-PEBBLE CONGLOMERATES



To test whether there are systematic diagenetic differences between QPC and polymict conglomerates, we examined conglomerates that had been subject to diagenetic processes for several hundred million years (Table 2; see footnote 1). Data were collected from interclast material because (1) diagenetic processes operate throughout the rock and so can be examined either at the clast or interclast level; (2) diagenetic products are concentrated in the interclast spaces; and (3) the mineralogic products of diagenesis are fine grained and therefore best examined microscopically.

All samples were unidirectional-flow deposits, clast supported, and well sorted. Sampled intervals contained traction structures such as parallel stratification, crude cross-stratification, imbrication, or intercalated flat-laminated sand lenses, indicating vigorous current activity. Such deposits contain very small amounts of primary matrix (Visher, 1969). Samples were not taken from diamictites, Bouma sequences, or other sediment gravity-flow deposits.

The interclast material in polymict conglomerates is dominated by sand-sized grains, with low proportions of matrix (Table 2 [see footnote 1]; Fig. 3). Water-laid QPC, however, consistently have large proportions of fine matrix. We interpret matrix volumes >10% (a conservative figure; see Visher, 1969) as secondary because they are inconsistent with the sample sedimentology. In addition, petrologic

**Figure 3.** Point-count data for interclast material in conglomerates. Value for pressure-solved grain contacts is sum of nested + interpenetrating contacts. Complete data are given in Table 2 (see text footnote 1).






**Figure 4. Pressure-solved grain contacts in quartz-pebble conglomerates from Mazatzal Group. Framework grains are quartz. Interstitial material is sericitic matrix.**

indicators, such as matrix-filled pore spaces and floating sand-sized grains (Dickinson, 1970), are commonly seen in the high-matrix rocks.

The most likely source of the matrix is the breakdown of labile detrital components (Dickinson, 1970). Postdepositional addition is unlikely because the observed volumes often greatly exceed the 3%–9% maximum volume reported for infiltrated material (Morales and De Ros, 1992). In addition, infiltrated clays, which are deposited from through-flowing muddy water, generally form grain coatings (Matlack et al., 1989), whereas the matrix in the QPC completely occludes pore space.

QPC exhibit substantially more pressure solution than do polymict conglomerates. Grains in polymict conglomerates generally have convex to straight grain edges, with tangential or linear contacts (Table 2 [see footnote 1]; Fig. 3). Grain contacts in QPC, in contrast, may be strongly modified (Fig. 4), with high proportions (15%–45%) of nested and interpenetrating contacts (definitions follow Taylor, 1950). The strong petrographic distinctions between QPC and polymict conglomerates are summarized in Figure 3. Polymict samples cluster at the low-matrix, low-pressure-solution corner of the diagram, whereas the QPC data spread out to much higher values of one or both parameters.



**Figure 5. Strength data for clasts from Tertiary Gila assemblage conglomerates of Tonto Basin.**

#### CONGLOMERATE COMPOSITIONS: SAME SOURCE AREA, DIFFERENT AGES

Conglomerates with very similar source rocks and depositional settings but very different ages (Proterozoic and Tertiary) are found in central Arizona. The Proterozoic Mazatzal Group is in depositional contact with a varied provenance, including silicified rhyolite (now chert), mafic igneous rocks, granitic plutons, schist, and quartzite (Wrucke and Conway, 1987), and the basal conglomerate was deposited in a proximal alluvial fan system (Trevena, 1979). The Miocene-Pliocene Gila Conglomerate was also deposited on an alluvial fan complex (Scarborough, 1989; Nations, 1990). Its source area includes the same rocks that contributed to the Mazatzal Group, in addition to Tertiary basalt and the Mazatzal Group. The clast compositions of the two units are, however, profoundly different. The Tertiary Gila Conglomerate contains quartzite, rhyolitic chert, schist, granite, and basalt; non-quartz clasts range from 22% to 55% by volume (Lang, 1999). Clasts in the Proterozoic Mazatzal Group conglomerate, in contrast, are 100% quartzose, consisting of rhyolitic chert, quartzite, and vein quartz (Trevena, 1979; Bayne, 1987; Cox and Lowe, 1995).

The Mazatzal Group has undergone extensive diagenetic alteration. Almost all interstitial material in the Mazatzal Group conglomerate has been converted to phyllosilicate secondary matrix, and the associated quartzites are diagenetic quartz arenites (Cox and Lowe, 1996). In addition, interstitial grain contacts in the conglomerates are strongly pressure solved (Fig. 4). We infer that the Mazatzal Group conglomerate originated as a polymict deposit similar to the Gila Conglomerate. The secondary matrix represents the diagenetic breakdown of labile clasts, and the pressure solution records dissolution, mass

transfer of soluble material, and consequent volume loss.

Processes operating currently provide a snapshot of the early stages of this transformation and suggest preadaptation to diagenetic QPC formation. The Tertiary Gila Conglomerate has never been buried. It is uncompacted and unlithified, but there has been substantial postdepositional compositional modification. Basalt, granite, and schist cobbles retain their shapes and textural characteristics and are therefore identifiable; but they have been altered to clays in situ and have little or no internal strength (Fig. 5). Clasts with a low strength index will not survive as recognizable lithologies during compaction and diagenesis, but will be crushed and redistributed as phyllosilicate-rich interstitial material. The high surface area of the fine-grained alteration products also makes them more susceptible to dissolution and mass transfer. We infer that much compositional modification leading to the development of diagenetic QPC may take place before deep burial.

#### CONCLUSIONS

This study indicates that, in many cases, QPC owe their composition to diagenetic processes. It is probable that clast disintegration and dissolution, followed by porosity collapse and pressure solution along grain boundaries, is responsible for the composition and texture of QPC. The volume of phyllosilicate-rich secondary material remaining in QPC is a minimum estimate of the volume of labile clasts lost during diagenesis, because a substantial proportion of the original clast material has probably been removed in solution. The extent of mass transfer is impossible to quantify, but the proportion of pressure-solution volume loss may provide a reasonable estimate of the relative magnitude of the effect.

We do not suggest that all QPC formed diagenetically. Intense weathering and protracted transport can certainly produce such deposits. However, the prevalence of QPC in settings where conditions are not optimal for their formation as primary deposits strongly suggests that postdepositional modification often plays a major role. The possibility that diagenesis may substantially alter conglomerate compositions means that careful examination of rock texture and interclast composition is required before paleoenvironmental conclusions can be drawn.

#### ACKNOWLEDGMENTS

This research was supported by National Science Foundation grant EAR-9814945 and American Chemical Society Petroleum Research Fund grant 34981-B8. We thank Clint Cowan and Marc Hendrix for very thoughtful and thorough reviews.

#### REFERENCES CITED

- Abbott, P.L., and Peterson, G.L., 1978, Effects of abrasion durability on conglomerate clast populations: Examples from Cretaceous and Eocene conglomerates of the San Diego area, California: *Journal of Sedimentary Petrology*, v. 48, p. 31–42.
- Abdel Wahab, A., 1998, Diagenetic history of Cambrian quartzarenites, Ras Dīb–Zeit Bay area, Gulf of Suez, Eastern Desert, Egypt: *Sedimentary Geology*, v. 121, p. 121–140.
- Bayne, B.J., 1987, Depositional analysis of conglomerates in the Mazatzal Group and related strata, central Arizona: Flagstaff, Northern Arizona University, 182 p.
- Boggs, S., 1992, *Petrology of sedimentary rocks*: New York, Macmillan Publishing Co., 575 p.
- Boggs, S.J., 2001, *Principles of sedimentology and stratigraphy*: Englewood Cliffs, New Jersey, Prentice-Hall, 726 p.
- Cavazza, W., and Gandolfi, G., 1992, Diagenetic processes along a basin-wide marker bed as a function of burial depth: *Journal of Sedimentary Petrology*, v. 62, p. 261–272.
- Cherichetti, L., Doolan, B., and Mehrtens, C., 1998, The Pinnacle Formation: A late Precambrian rift valley fill with implications for Iapetus rift basin evolution: *Northeastern Geology and Environmental Sciences*, v. 20, p. 175–185.
- Cox, R., and Lowe, D.R., 1995, Compositional evolution of coarse clastic sediments in the southwestern United States from 1.8–0.2 Ga and implications for relationships between the development of crustal blocks and their sedimentary cover: *Journal of Sedimentary Research*, v. A65, p. 477–494.
- Cox, R., and Lowe, D.R., 1996, The effects of secondary matrix on detrital modes of sandstones, and the relationships between sandstone chemistry and tectonic setting: Implications for provenance studies: *Journal of Sedimentary Research*, v. 66, p. 548–558.
- Cronin, B.T., and Kidd, R.B., 1998, Heterogeneity and lithotype distribution in ancient deep-sea canyons; Point Lobos deep-sea canyon as a reservoir analogue: *Sedimentary Geology*, v. 115, p. 315–349.
- Dal Cin, R., 1968, Climatic significance of roundness and percentage of quartz in conglomerates: *Journal of Sedimentary Petrology*, v. 38, p. 1094–1099.
- Dickinson, W.R., 1970, Interpreting detrital modes of graywacke and arkose: *Journal of Sedimentary Petrology*, v. 40, p. 695–707.
- Ethridge, F.G., Tyler, N., and Burns, L.K., 1984, Sedimentology of a Precambrian quartz-pebble conglomerate, southwest Colorado, in Koster, E.H., and Steel, R.J., eds., *Sedimentology of gravels and conglomerates: Canadian Society of Petroleum Geologists Memoir 10*, p. 165–174.
- Garrels, R.M., Mackenzie, F.T., and Siever, R., 1972, Sedimentary cycling in relation to the history of the continents and oceans, in Robertson, E.C., ed., *The nature of the solid earth*: New York, McGraw-Hill, p. 93–121.
- Gilluly, J., 1969, Geologic perspective and the completeness of the geologic record: *Geological Society of America Bulletin*, v. 80, p. 2303–2311.
- Glover, G., 1982, A study of the Bend Conglomerate in SE Maryetta area, Boonesville Field, Jack County, Texas, in Martin, C.A., ed., *Petroleum geology of the Fort Worth Basin and Bend Arch area*: Dallas, Texas, Dallas Geological Society, p. 353–364.
- Gregor, C.B., 1985, The mass-age distribution of Phanerozoic sediments, in Snelling, N.J., ed., *The chronology of the geologic record*: Geological Society [London] Memoir 10, p. 284–289.
- Harris, N.B., 1989, Diagenetic quartzarenite and destruction of secondary porosity: An example from the Middle Jurassic Brent sandstone of northwest Europe: *Geology*, v. 17, p. 361–364.
- Houseknecht, D.W., 1987, Assessing the relative importance of compaction processes and cementation to reduction of porosity in sandstones: *American Association of Petroleum Geologists Bulletin*, v. 71, p. 633–642.
- Houseknecht, D.W., 1989, Assessing the relative importance of compaction processes and cementation to reduction of porosity in sandstones: Reply: *American Association of Petroleum Geologists Bulletin*, v. 73, p. 1277–1279.
- Kingsley, C.S., 1984, Dagbreek fan-delta: An alluvial placer to prodelta sequence in the Proterozoic Welkom goldfield, Witwatersrand, South Africa, in Koster, E.H., and Steel, R.J., eds., *Sedimentology of gravels and conglomerates*: Canadian Society of Petroleum Geologists Memoir 10, p. 321–330.
- Kraus, M.J., 1984, Sedimentology and tectonic setting of early Tertiary quartzite conglomerates, northwest Wyoming, in Koster, E.H., and Steel, R.J., eds., *Sedimentology of gravels and conglomerates*: Canadian Society of Petroleum Geologists Memoir 10, p. 203–216.
- Lang, N., 1999, Correlation and mapping of a Tertiary alluvial fan complex, central Arizona: *Proceedings of the Keck Research Symposium in Geology*, v. 12, p. 247–250.
- Matlack, K.S., Houseknecht, D.W., and Applin, K.R., 1989, Emplacement of sand into clay by infiltration: *Journal of Sedimentary Petrology*, v. 59, p. 77–87.
- Milliken, K.L., 1988, Loss of provenance information through subsurface diagenesis in Plio-Pleistocene sandstones, northern Gulf of Mexico: *Journal of Sedimentary Petrology*, v. 58, p. 992–1002.
- Milliken, K.L., Mack, L.E., and Land, L.S., 1994, Elemental mobility in sandstones during burial: Whole-rock chemical and isotopic data, Frio Formation, south Texas: *Journal of Sedimentary Petrology*, v. A64, p. 788–796.
- Moraes, M.A.S., and De Ros, L.F., 1992, Depositional, infiltrated and authigenic clays in fluvial sandstones of the Jurassic Sergi Formation, Recôncavo Basin, northeastern Brazil, in Houseknecht, D.W., and Pittman, E.D., eds., *Origin, diagenesis and petrophysics of clay minerals in sandstones*: SEPM (Society for Sedimentary Geology) Special Publication 47, p. 197–208.
- Mosher, S., 1976, Pressure solution as a deformation mechanism in Pennsylvanian conglomerates from Rhode Island: *Journal of Geology*, v. 84, p. 355–364.
- Mosher, S., 1981, Pressure solution deformation of the Purgatory Conglomerate from Rhode Island: *Journal of Geology*, v. 89, p. 37–55.
- Mosher, S., Murray, D.P., Hermes, O.D., Grömet, L.P., and Hepburn, J.C., 1993, Alleghanian and Avalonian tectonism in southeastern New England, in Cheney, J.T., and Hepburn, J.C., eds., *Field trip guidebook for the northeastern United States*, 1993 Boston GSA, v. 2: Amherst, Massachusetts, University of Massachusetts, p. BB1–BB30.
- Nations, J.D., 1990, Late Cenozoic stratigraphy and tectonics of the Tonto Basin, central Arizona, in Gehrels, G.E., and Spencer, J.E., eds., *Geologic excursions through the Sonoran Desert region, Arizona and Sonora*: Arizona Bureau of Geology and Mineral Technology Special Paper 7, p. 24–30.
- Prothero, D.R., and Schwab, F., 1996, *Sedimentary geology*: New York, W.H. Freeman and Co., 575 p.
- Reimer, T.O., and Mossman, D.J., 1990, The Witwatersrand controversy revisited: *Economic Geology*, v. 85, p. 337–343.
- Robinson, A., and Spooner, E., 1984, Postdepositional modification of uraninite-bearing quartz-pebble conglomerates from the Quirke ore zone, Elliot Lake, Ontario: *Economic Geology*, v. 79, p. 297–321.
- Ronov, A.B., 1983, The Earth's sedimentary shell: Quantitative patterns of its structure, composition and evolution: *American Geological Institute Reprint Series* no. 5, 80 p.
- Ronov, A.B., Khain, V.E., Balukhovskiy, A.N., and Sesslavinsky, K.B., 1980, Quantitative analysis of Phanerozoic sedimentation: *Sedimentary Geology*, v. 52, p. 311–325.
- Scarborough, R., 1989, Cenozoic erosion and sedimentation in Arizona, in Jenney, J.P., and Reynolds, S.J., eds., *Geologic evolution of Arizona*: Arizona Geological Society Digest 17, p. 515–537.
- Selley, R.C., 2000, *Applied sedimentology*: New York, Academic Press, 521 p.
- Siebert, R.M., Moncure, G.K., and Lahann, R.W., 1984, A theory of framework grain dissolution in sandstones, in McDonald, D.A., and Surdam, R.C., eds., *Clastic diagenesis*: American Association of Petroleum Geologists Memoir 37, p. 163–175.
- Smith, N.D., 1967, A stratigraphic and sedimentologic analysis of some Lower and Middle Silurian clastic rocks of the north-central Appalachians [Ph.D. thesis]: Providence, Rhode Island, Brown University, 195 p.
- Surdam, R.C., Boese, S.W., and Crossey, L.W., 1984, The chemistry of secondary porosity, in McDonald, D.A., and Surdam, R.C., eds., *Clastic diagenesis*: American Association of Petroleum Geologists Memoir 37, p. 127–134.
- Taylor, J.M., 1950, Pore-space reduction in sandstone: *American Association of Petroleum Geologists Bulletin*, v. 34, p. 701–706.
- Trevena, A.S., 1979, Studies in sandstone petrology: Origin of the Precambrian Mazatzal Quartzite and provenance of detrital feldspar [Ph.D. thesis]: Salt Lake City, University of Utah, 390 p.
- Visher, G.S., 1969, Grain size distributions and depositional processes: *Journal of Sedimentary Petrology*, v. 39, p. 1074–1106.
- Wilkinson, M., Darby, D., Haszeldine, R.S., and Couples, G.D., 1997, Secondary porosity generation during deep burial associated with overpressure leak-off: Fulmar formation, United Kingdom Central Graben: *American Association of Petroleum Geologists Bulletin*, v. 81, p. 803–813.
- Wrucke, C.T., and Conway, C.M., 1987, Geologic map of the Mazatzal wilderness and contiguous roadless area, Gila, Maricopa and Yavapai Counties, Arizona: U.S. Geological Survey Open-File Report 87-664, 22 p.
- Youngson, J.H., and Craw, D., 1996, Recycling and chemical mobility of alluvial gold in Tertiary and Quaternary sediments, central and East Otago, New Zealand: *New Zealand Journal of Geology and Geophysics*, v. 39, p. 493–508.

Manuscript received September 4, 2001  
 Revised manuscript received November 13, 2001  
 Manuscript accepted December 14, 2001

Printed in USA